



Contribution ID: 364

Type: **not specified**

P4.028 Development of cesium-free negative hydrogen/deuterium ion source by sheet plasma

Thursday, 20 September 2018 11:00 (2 hours)

Negative ions play an essential role for Neutral Beam Injection (NBI) system of steady state magnetic nuclear fusion. We have development of negative ion source in a cesium-free discharge by the magnetized sheet plasma device, TPD-Sheet IV [1]. Negative ions are formed by volume-production, that is, the dissociative attachment of low energy electrons ($T_e = 1-2$ eV) to highly vibrationally excited molecules. These molecules are attributed to the electron-impact excitation of molecules by high energy electrons ($T_e > 10$ eV) in the plasma column. Negative hydrogen/deuterium ion beams are extracted through the small hole (6mm in diameter) with an extraction voltage VEX of 3 kV at a neutral gas pressure P_s of 0.3 Pa. The maximum negative hydrogen ion beam current density J_{H^-} is about 7.0 mA/cm² at discharge current I_d of 50 A. On the other hand, the negative deuterium ion beam current density J_{D^-} is obtained about 4.0 mA/cm² at I_d of 80 A.

This study is supported by the LHD Joint Project of the National Institute for Fusion Science (NIFS16KOAR021).

Presenter: HANAI, Keito (Science Tokai University Kitakaname)

Session Classification: P4